



(11) Publication number : **0 307 128 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication of patent specification :
08.04.92 Bulletin 92/15

(51) Int. Cl.⁵ : **A61K 31/71, C07H 17/00**

(21) Application number : **88308024.4**

(22) Date of filing : **31.08.88**

(54) **Azithromycin and derivatives as antiprotozoal agents.**

The file contains technical information
submitted after the application was filed and
not included in this specification

(30) Priority : **10.09.87 PCT/US87/02317**

(43) Date of publication of application :
15.03.89 Bulletin 89/11

(45) Publication of the grant of the patent :
08.04.92 Bulletin 92/15

(84) Designated Contracting States :
AT BE CH DE ES FR GB GR IT LI LU NL SE

(56) References cited :
EP-A- 0 101 186
EP-A- 0 109 253
ANTIMICROBIAL AGENTS AND
CHEMOTHERAPY, vol. 32, no. 5, May 1988,
pages 755-757, American Society for Micro-
biology; F.G. ARAUJO et al.: "Azithromycin, a
macrolide antibiotic with potent activity
against toxoplasma gondii"
ANTIMICROBIAL AGENTS AND
CHEMOTHERAPY, vol. 32, no. 4, April 1988,
Pages 524-529, American Society for Micro-
biology; H.R. CHANG et al.: "In vitro effects of
four macrolides (Roxithromycin, Spiramycin,
Azithromycin [CP-62,993], and A-56268) on
toxoplasma gondii"
AIDS, vol. 2, suppl 1, 1988, pages S177-S182,
Gower Academic Journals Ltd; A. PITCHENIK:
"The treatment and prevention of mycobacteri-
al disease in patients with HIV infection"
ANTIMICROBIAL AGENTS AND
CHEMOTHERAPY, vol. 31, no. 2, February
1987, pages 345-348, American Society for
Microbiology; J.M. HOFFLIN et al.: "In vivo
synergism of Roxithromycin (RU 965) and in-
terferon against Toxoplasma gondii"

(56) References cited :
ANTIMICROBIAL AGENTS AND
CHEMOTHERAPY, vol. 31, no. 5, May 1987,
pages 811-812, American Society for Micro-
biology; M. WALSH et al.: "In viro evaluation of
CP-62,993, Erythromycin, Clindamycin, and
Tetracycline against Chlamydia trachomatis"
EUR: J. CLIN. MICROBIOL., vol. 6, no. 1, Feb-
ruary 1987, pages 1-2; F. ARAUJO et al.:
"Toxoplasmosis in immunocompromised
patients"
Antimicrobial Agents and Chemotherapy,
1988, 32(5), 755-757
**Journal of Chemotherapy of Infectious Dis-
eases and Malignancies**, 1989, Suppl.1

(73) Proprietor : **PFIZER INC.**
235 East 42nd Street
New York, N.Y. 10017 (US)

(72) Inventor : **Remington, Jack Samuel**
1995 Val Paraiso Avenue
Menlo Park California (US)

(74) Representative : **Moore, James William, Dr.**
Pfizer Limited Ramsgate Road
Sandwich Kent CT13 9NJ (GB)

EP 0 307 128 B1

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

1

5

10

25

35



45

50

(Ic) $R^1 = NH_2$, $R^2 = H$ 4"-amino-4"-deoxy-azithromycin
 (Id) $R^1 = H$, $R^2 = NH_2$ 4"-epi-4"-amino-4"-deoxy-azithromycin

5 Summary of the Invention

We have now found that the compounds of the formula (I), wherein one of R^1 and R^2 is hydrogen and the other is hydroxy or amino (conveniently named herein as azithromycin derivatives, *vide supra*) possess remarkably potent activity against *Toxoplasma* species, and so are valuable in pharmaceutical compositions for a method of treating or preventing protozoal infections in mammals, including man. These compounds are especially valuable in the treatment of toxoplasmosis, an infection due to a strain of *Toxoplasma gondii*, which, as noted above, is a particular problem in pregnant women and in immune compromised patients.

Detailed Description of the Invention

15 The present invention is readily carried out. The compounds of the formula (I) are prepared according to the methods of U.S. Patents 4,474,768, 4,512,982 and 4,526,889, cited above. A particularly valuable form of azithromycin (Ia) for this purpose is azithromycin dihydrate prepared according to methods disclosed in Examples below.

20 The utility of the compounds of the formula (I) in the treatment or prevention of protozoal infections in mammals is demonstrated by their remarkable activity in model *Toxoplasma gondii* infections in mice. For example, we have found azithromycin (Ia) to have potent *in vivo* activity against murine toxoplasmosis. Mice infected intraperitoneally with 10^2 tachyzoites of the virulent RH strain of *T. gondii* and treated 24 hours later with 200 mg azithromycin/kg/day orally by gavage (solubilized in polyethylene glycol 200) for 10 days all survived. Concentrations of 100 or 50 mg/kg resulted in 80 and 20% survival, respectively. Further experiments revealed that one daily dose of 200 mg/kg for each of 3 days after infection resulted in 100% survival of mice infected with 10^3 RH tachyzoites. Moreover, this concentration of the drug protected 100% of infected mice when administered as late as 72 hours after infection with 10^2 RH tachyzoites. Additional experiments revealed that 70% of mice infected intracerebrally with 10^4 tachyzoites of the C56 strain of *T. gondii* and treated with 200 mg/kg/day for 10 days survived, but only 10% of untreated controls survived. These results indicate that azithromycin is highly effective against infection with *T. gondii*. (See Hofflin *et al.*, cited above, and references there cited, for more detailed descriptions of these murine toxoplasmosis models).

35 In the treatment or prevention of systemic protozoal infections in mammals, particularly toxoplasmosis in man due to strains of *Toxoplasma gondii*, the compounds of the formula (I), including the pharmaceutically acceptable salts thereof, are dosed orally or parenterally. Oral dosage will generally be preferred, particularly in cases where the drug is dosed chronically as a preventive measure. However, particularly in acute administration for severe cases of toxoplasmosis, parenteral administration may be preferred, a matter to be determined at the discretion of the attending physician. The preferred dosage range is 5-100 mg per kg of body weight per day, in single or divided daily doses, regardless of the route of administration. In special situations, particularly in life-threatening cases of infection, higher doses may be prescribed at the discretion of the attending physician.

40 When used to treat or prevent a systemic protozoal infection in a mammal, particularly toxoplasmosis in man, the compounds of the formula (I), including the pharmaceutically acceptable salts thereof, can be dosed alone, but are preferably dosed in the form of pharmaceutical compositions comprising the active compound and a pharmaceutically-acceptable carrier or diluent. Such pharmaceutical compositions, whether for oral or parenteral administration, are prepared according to conventional methods of pharmacy, for example, as disclosed in U.S. Patents 4,474,768, 4,512,982 and 4,526,889, cited above.

The present invention is illustrated by the following example.

50 EXAMPLE 1

Non-Hygroscopic Azithromycin Dihydrate

Method A

55 The hygroscopic monohydrate of Preparation 1 (100 g; water-content:3.1%), tetrahydrofuran (220 ml) and diatomaceous earth (5 g) were combined in a 500 ml Erlenmeyer flask, stirred for 30 minutes and filtered with 20 ml of tetrahydrofuran wash. The combined filtrate and wash was transferred to a 3 liter round bottom flask.

The solution was stirred vigorously and H₂O (2.0 ml) was added. After 5 minutes, hexane (1800 ml) was added over 5 minutes, with continued vigorous stirring. Following an 18 hour granulation period, title product was recovered by filtration with 1 x 10 ml hexane wash, and dried in vacuo to 4.6±0.2% H₂O by Karl Fischer, 89.5 g.

5

Method B

The hygroscopic monohydrate of Preparation 1 (197.6 g) and tetrahydrofuran (430 ml) were charged to a reactor and the mixture stirred to achieve a milky white solution. Activated carbon (10 g) and diatomaceous earth (10 g) were added and the mixture stirred for 15 minutes, then diluted with 800 ml of hexane and filtered with suction over a pad of diatomaceous earth with 250 ml of hexane for wash. The combined filtrate and wash was diluted to 2500 ml with hexane and warmed to 34°C. With stirring, 24.7 ml of H₂O was added. The mixture was allowed to cool to room temperature, granulated for five hours and title product recovered and dried as in Method A, 177.8 g.

The dihydrate melts sharply at 126°C (hot stage, 10°/minute); differential scanning calorimetry (heating rate, 20°C/minute) shows an endotherm at 127°C; thermal gravimetric analysis (heating rate 30°C/minute) shows a 1.8% weight loss at 100°C and a 4.3% weight loss at 150°C; ir (KBr) 3953, 3553, 3488, 2968, 2930, 2888, 2872, 2827, 2780, 2089, 1722, 1664, 1468, 1426, 1380, 1359, 1344, 1326, 1318, 1282, 1270, 1252, 1187, 1167, 1157, 1123, 1107, 1082, 1050, 1004, 993, 977, 955, 930, 902, 986, 879, 864, 833, 803, 794, 775, 756, 729, 694, 671, 661, 637, 598, 571, 526, 495, 459, 399, 374, 321 and 207 cm⁻¹; [α]_D²⁵ = -41.4° (c=1, CHCl₃).

20

Anal. Calcd. for C₃₈H₇₂N₂O₁₂·2H₂O:

C, 58.14; H, 9.77; N, 3.57; OCH₃, 3.95; H₂O, 4.59.

Found:

25

C, 58.62; H, 9.66; N, 3.56; OCH₃, 4.11; H₂O, 4.49.

Neutralization Equivalent (0.5N HCl in 1:1 CH₃CN:H₂O):

Calcd.: 374.5. Found: 393.4.

30

Samples of a dihydrate, slightly over dried to contain 4.1% water (less than theoretical) rapidly picked-up water at 33%, 75% or 100% relative humidities to achieve the theoretical water content (4.6%) for the dihydrate. At 33% and 75% relative humidities, water content remained essentially constant for at least 4 days. At 100% relative humidity, the water content further rose to about 5.2, where it remained essentially constant of the next three days.

A sample of the same dihydrate, maintained at 18% relative humidity gradually lost water. At four days, the water content was 2.5% and at 12 days, 1.1%.

35

EXAMPLE 2

Azithromycin Powder for Oral Suspension

40

The following powdered ingredients were thoroughly blended:

45

50

55

	Azithromycin Dihydrate (1200 g on anhydrous basis)	1268.5 g
5	Sucrose	23000 g
	Sodium phosphate tribasic dodecahydrate	250 g
10	Sodium benzoate	90 g
	Hydroxypropylcellulose	40 g
	Xanthan gum	40 g
15	Certified food coloring agent(s) in solid form	3 g or as required to achieve the desired color
20	Fruit and/or vanilla flavoring agents in solid form	440 g or as required to achieve the desired taste

25 The resulting blend contains 47.75 mg of azithromycin activity per gram. Amber screw cap bottles (60 ml) are filled with 10.47 g of the blend. Prior to oral administration as a suspension, distilled water is added (25 ml) and the mixture shaken. One teaspoon (5 cc) of this mixture provides a 100 mg dose of azithromycin. Higher or lower doses are achieved by appropriate modification of the dosage volume.

30 EXAMPLE 3

Azithromycin Capsules (250 mg) for Oral Administration

35 The following ingredients were accurately weighed, combined, and blended in a suitable blender for 15 minutes.

	Hydrated azithromycin	3360.9 g*
40	*(3250.0 g on an anhydrous basis)	
	Anhydrous lactose	2015.9 g
	Corn starch	611.0 g

45 The blended material was milled through Fitz JT mill with a No. 2A plate (0.093") at slow speed with knives forward, the milled mixture blended for an additional 15 minutes, and weighed. The resulting milled and blended mixture (5977.2 g) was then blended for 5 minutes with a 9:1 lubricant mixture of magnesium stearate:sodium lauryl sulfate (91.65 g), the further blend slugged on a Stokes DD-2 fitted with six stations of 3/4" flat faced punches, and the slugs granulated by remilling and additionally blending as specified above. Additional 9:1 lubricant (29.5 g) was blended with the resulting granulated blend (5869 g) and the material encapsulated into #0 capsules on a Zanasi RM-63 capsule machine at a fill weight of 483±23 mg to yield capsules containing no more than 275 mg and no less than the desired 250 mg of azithromycin activity.

By appropriately modifying the capsule size, the fill weight and the proportion of azithromycin in the blend, capsules containing 100 mg, 125 mg, 375 mg or 500 mg of azithromycin activity are prepared.

55 4"-Epi-azithromycin, 4"-amino-4"-deoxy-azithromycin and 4"-epi-4"-amino-4"-deoxy-azithromycin capsules are prepared in like manner, substituting in equal weight of the active ingredient (corrected for potency as free base) for azithromycin.

EXAMPLE 4Azithromycin Tablets (250 Mg) for Oral Administration

5 The following ingredients were accurately weighed, combined and blended in a suitable blender for 30 minutes:

	Azithromycin dihydrate	14245.0 g*
10	*(13,485.0 g on an anhydrous basis)	
	Dibasic calcium phosphate	22205.0 g
	AC-DI-SOL ^(R)	1620.0 g
15	Magnesium stearate	1242.7 g

20 The blend was milled in a Fitzpatrick D comminutor fitted with a No. 3 plate (0.125") with knives forward at 3600 rpm, then blended for an additional 30 minutes. To the resulting milled blend (39,192 g) was added an additional 783.8 g of magnesium stearate and blending continued for 5 minutes. The mixture was then slugged according to the preceding example, and remilled as immediately above, and blended for 5 minutes. Additional magnesium stearate (394.5 g) was added to the resulting granulated blend (39,445 g), blending was continued for 5 minutes, and the mixture tableted on a Killian tableting machine with forced feeder and 32" x 5/8" upper and lower oval shaped punches, each tablet having a weight of 787 mg \pm 37 mg, each containing no less than 250 mg and no more than 275 mg of azithromycin activity.

EXAMPLE 5Azithromycin for I.V. or I.M. Injection

30 In a sterile environment and using sterile, particle free equipment and components, 10,949 g of water for injection was placed in a compounding flask. Anhydrous citric acid, 494.4 g was added and dissolved with agitation. In a separate flask 310 g of sodium hydroxide was dissolved in 690 g of water. A portion of the latter (755 g) was used to adjust the pH of the citric acid from 1.63 to 5.09 ± 0.02 . Azithromycin dihydrate 670.0 g (equivalent to 642.5 g of anhydrous base) was added, and the mixture adjusted to pH 6.60 ± 0.1 with 4.0 g additional of the sodium hydroxide solution. Water (6076.5 g) was added to bring the resulting solution to a final weight of 18,948.9 g. If desired, the solution is sterile filtered at this stage, using a millipore filter. Using a filling machine, 50 ml flint type vials were each filled with 15.06 ± 0.45 g of this solution, loosely stoppered with gray teflon stoppers, and freeze dried to yield stoppered vials each containing 51 ± 1.5 mg of azithromycin activity in the form of freeze dried solids. Prior to i.m. or i.v. injection, water for injection (10 ml) is added by injection by syringe through the stopper, and the freeze dried solids redissolved by shaking. Virtually the entire contents of the vial is taken up into the syringe and injected either i.v. or i.m.

PREPARATION 1Hygroscopic Azithromycin Monohydrate

45 Substantially following the methylation procedure of Kobrehel et al., U.S. Patent 4,517,359; and the crystallization procedure of Bright, U.S. Patent 4,474,768; 9-deoxy-9a-aza-9a-homoerythromycin A (previously called 11-aza-10-deoxy-10-dihydroerythromycin A; 100 g, 0.218 mol) was dissolved with stirring in 400 ml CHCl_3 . Formic acid (98%; 10.4 ml, 0.436 mol) and formaldehyde (37%; 16.4 ml, 0.349 mol) were added over 4-5 minutes, and the mixture heated at reflux for 20 hours. The mixture was cooled to ambient temperature, diluted with 400 ml H_2O and adjusted to pH 10.5 with 50% NaOH. The aqueous layer was separated and extracted 2 x 100 ml with fresh CHCl_3 . The organic layers were combined, stripped *in vacuo* to 350 ml, twice diluted with 50 ml of ethanol and reextracted to 350 ml, and finally diluted with 1000 ml H_2O over a 1 hour period, pausing for 15 minutes as a slurry began to develop after the addition of about 250 ml of H_2O . Title product was recovered by filtration and dried in air at 50°C for 24 hours, 85 g; mp 136°C; differential thermal analysis (heating rate 20°C/minute) shows an endotherm at 142°C; thermal gravimetric analysis (heating rate 30°C/minute) shows a

2.6% weight loss at 100°C and a 4.5% weight loss at 150°C; water content 3.92%; ethanol content 1.09%.

Anal. Calcd. for $C_{38}H_{72}N_2O_{12}$ (corrected for ethanol and water content):

C, 58.46; H, 9.78; N, 3.74; Alkoxy, 4.67.

Found: C, 58.40; H, 9.29; N, 3.50; Alkoxy, 4.52.

- 5 A sample of the monohydrate (having a water content of 3.2%) was maintained at 18% relative humidity for 14 days. The sample lost water over the first 24 hours to yield monohydrate having the theoretical water content (2.35%). The water content then remained substantially constant over 14 days, a value of 2.26% being recorded at 14 days.

- 10 At 33% relative humidity the water content of a sample of the same monohydrate rapidly rose to 5.6% where it remained substantially steady for at least three days. Similarly at 75% and 100% relative humidity, the water content rose rapidly, but was now maintained at even higher levels, 6.6% and 7.2%, respectively, for at least 3 days.

15 Claims

1. Use of an anti-Toxoplasma gondii effective amount of:

azithromycin;

4"-epi-azithromycin;

- 20 4"-amino-4"-deoxy-azithromycin; or

4"-epi-4"-amino-4"-deoxy-azithromycin;

or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use in the treatment or prevention of an infection in a mammal due to a strain of Toxoplasma gondii.

- 25 2. The use of a compound according to claim 1 wherein said compound is azithromycin or a pharmaceutically acceptable salt thereof.

3. The use of a compound according to claim 1 wherein said compound is 4"-epi-azithromycin or a pharmaceutically acceptable salt thereof.

4. The use of a compound according to claim 1 wherein said compound is 4"-amino-4"-deoxy-azithromycin or a pharmaceutically acceptable salt thereof.

- 30 5. The use of a compound according to claim 1 wherein said compound is 4"-epi-4"-amino-4"-deoxy-azithromycin or a pharmaceutically acceptable salt thereof.

Patentansprüche

35

1. Verwendung einer gegen Toxoplasma gondii wirksamen Menge von Azithromycin,

4"-Epi-azithromycin,

4"-Amino-4"-desoxy-azithromycin oder

4"-Epi-4"-amino-4"-desoxy-azithromycin

- 40 oder eines pharmazeutisch annehmbaren Salzes hiervon für die Herstellung eines Medikaments zur Verwendung bei der Behandlung oder Verhütung einer Infektion eines Säugers durch einen Stamm von Toxoplasma gondii.

2. Verwendung einer Verbindung nach Anspruch 1, wobei die Verbindung Azithromycin oder ein pharmazeutisch annehmbares Salz hiervon ist.

- 45 3. Verwendung einer Verbindung nach Anspruch 1, wobei die Verbindung 4"-Epi-azithromycin oder ein pharmazeutisch annehmbares Salz hiervon ist.

4. Verwendung einer Verbindung nach Anspruch 1, wobei die Verbindung 4"-Amino-4"-desoxy-azithromycin oder ein pharmazeutisch annehmbares Salz hiervon ist.

- 50 5. Verwendung einer Verbindung nach Anspruch 1, wobei die Verbindung 4"-Epi-4"-amino-4"-desoxy-azithromycin oder ein pharmazeutisch annehmbares Salz hiervon ist.

Revendications

- 55 1. Utilisation d'une quantité efficace contre Toxoplasma gondii de :

l'azithromycine ;

la 4"-épi-azithromycine ;

la 4"-amino-4"-dé xy-azithromycine ; u

la 4"-épi-4"-amino-4"-déoxy-azithromycine ;

ou d'un sel acceptable du point de vue pharmaceutique de ce composé, pour la préparation d'un médicament destiné à être utilisé dans le traitement curatif ou préventif d'une infection due à une souche de Toxoplasma gondii chez un mammifère.

5 2. Utilisation d'un composé suivant la revendication 1, dans laquelle ledit composé est l'azithromycine ou un sel pharmaceutiquement acceptable de ce composé.

3. Utilisation d'un composé suivant la revendication 1, dans laquelle le composé est la 4"-épi-azithromycine ou un sel pharmaceutiquement acceptable de ce composé.

10 4. Utilisation d'un composé suivant la revendication 1, dans laquelle le composé est la 4"-amino-4"-déoxy-azithromycine ou un sel pharmaceutiquement acceptable de ce composé.

5. Utilisation d'un composé suivant la revendication 1, dans laquelle le composé est la 4"-épi-4"-amino-4"-déoxy-azithromycine ou un sel pharmaceutiquement acceptable de ce composé.

15

20

25

30

35

40

45

50

55